

**PIOJOS**

# los Hechos de los Piojos


Los niños en la escuela preescolar y primaria son los más propensos a contraer piojos – pequeños insectos parásitos sin alas que se alimentan de cantidades pequeñas de sangre extraídas del cuero cabelludo de los humanos.

**COMPARACIÓN DE TOMAÑO**



<p><b>1</b></p> <p><b>LIENDRE</b></p> <p>Se ponen huevos diminutos en la base del tallo del cabello y a menudo se confunden con caspa.</p> 	<p><b>2</b></p> <p><b>NINFA</b></p> <p>Las liendres eclosionan en ninfas y se vuelven de color marrón rojizo después de alimentarse de sangre humana.</p> 	<p><b>3</b></p> <p><b>ADULTO</b></p> <p>Los piojos hembras pueden poner hasta 100 liendres durante sus 30-45 días de vida.</p> 
--	---	---

<p><b>1</b></p> <p>UNA SEMANA</p>	<p><b>2</b></p> <p>UNA SEMANA</p>	<p><b>3</b></p> <p>HASTA 30 DÍAS</p>
-----------------------------------	-----------------------------------	--------------------------------------



[www.LiceRemovalNOVA.com](http://www.LiceRemovalNOVA.com)  
(703) 303-1576

Los piojos no brincan ni vuelan. El contacto de **CABEZA A CABEZA** es la manera única que propaganan los piojos.



Los piojos casi nunca propaganan por medio de **SOMBREROS, CEPILLOS** u otros artículos.



**LAVE Y SEQUE** la ropa de cama que haya usado en las últimas 24 horas cualquier miembro de la familia que esté bajo tratamiento.



Ayude a prevenir la propagación de piojos usando el cabello en un **MOÑO APRETADO O TRENZAS**.



Los piojos **NO VIVEN** en sus mascotas, cuerpo o juguetes de peluche.




**HEAD LICE**

# the Facts of Lice


Children in preschool and elementary school are the most likely to get head lice - tiny, wingless parasitic insects that feed on tiny amounts of blood drawn from the scalp of humans.

Lice cannot jump or fly. The only way for them to spread is **HEAD-TO-HEAD CONTACT.**



Head lice almost never spread by **HATS, BRUSHES,** or other items.

**WASH AND DRY** bedding used in the past 24 hours with the hottest settings.




Help prevent the spread of lice by wearing hair in a tight **BUN** or **BRAIDS.**


Head lice **DO NOT LIVE ON** your pets, body, or stuffed toys.



**SIZE COMPARISON**



<p><b>1</b></p> <p><b>NIT</b></p> <p>Tiny eggs are laid at base of hair shaft and are often confused with dandruff.</p> 	<p><b>2</b></p> <p><b>NYMPH</b></p> <p>Nits hatch into baby insects and become reddish-brown after feeding on human blood.</p> 	<p><b>3</b></p> <p><b>ADULT</b></p> <p>Females can lay up to 100 eggs in their 30 to 45-day life span.</p> 
---	--	---



LiceRemovalNOVA.com  
(703) 303-1576